arXiv:1211.3538v1 [quant-ph] 15 Nov 2012

The Schmidt modes of biphoton qutrits: Poincaré-sphere representation

M V Chekhova[1,2] **and M V Fedorov**[3,4]

[1]Department of Physics, M V Lomonosov Moscow State University, Moscow, Russia
[2]Max-Planck-Institute for the Science of Light, Erlangen, Germany
[3]A M Prokhorov General Physics Institute, Russian Academy of Science, Moscow, Russia.
[4]Moscow Institute of Physics and Technology, Dolgoprudny, Moscow Region, Russia

Abstract. For a general-form polarization biphoton qutrit, physically corresponding to a pair of arbitrarily polarized photons in a single frequency and wavevector mode, we explicitly find polarization Schmidt modes. A simple method is suggested for factorizing the state vector and the explicit expressions for the factorizing photon creation operators are found. The degrees of entanglement and polarization of a qutrit are shown to depend directly on the commutation features of the factorizing operators. Clear graphic representations for the Stokes vectors of the qutrit state as a whole, its Schmidt modes, and factorizing single-photon creation operators are given, based on the Poincaré sphere.

PACS numbers: 03.67.Bg, 03.67.Mn, 42.65.Lm

1. Introduction

Biphoton qutrits are the simplest two-photon formations finding applications in the science of quantum information. In polarization biphoton qutrits two photons are assumed to have only one degree of freedom, polarization, with identical spatial and spectral features. Such states are usually produced via collinear spontaneous parametric down-conversion (SPDC) degenerate in the frequencies of the photons. The general form of a biphoton qutrit state vector is given by the superposition

$$|\Psi\rangle = C_1|2_H\rangle + C_2|1_H,\,1_V\rangle + C_3|2_V\rangle, \tag{1.1}$$

where $|2_H\rangle = \frac{a_H^{\dagger 2}}{\sqrt{2}}|0\rangle$ and $|2_V\rangle = \frac{a_V^{\dagger 2}}{\sqrt{2}}|0\rangle$ are, correspondingly, biphoton state vectors with both photons having either horizontal or vertical polarization, whereas $|1_H,\,1_V\rangle = a_H^\dagger a_V^\dagger|0\rangle$ is the state vector of two photons with different polarization, horizontal and vertical, $a_H^\dagger$ and $a_V^\dagger$ are creation operators for photons with horizontal and vertical polarizations. The constants $C_{1,2,3}$ obey the normalization condition $|C_1|^2 + |C_2|^2 + |C_3|^2 = 1$. In the general case $C_{1,2,3}$ can be complex. But as the phase of the state $|\Psi\rangle$ as a whole does not affect any measurable characteristics of qutrits, it can be chosen, e.g., in a way making C_1 real and positive with the parameters C_2 and C_3 being complex (if $C_1 = 0$ any other of two remaining parameters C_2 or C_3 can be taken real and positive). In the general case this leaves four free parameters determining states of qutrits, e.g., $|C_2|$ and $|C_3|$ plus their phases φ_2 and φ_3.

The main two physically important characteristics of biphoton qutrits are their entanglement and polarization. Polarization features of biphoton qutrits have been widely investigated since 1999 [1, 2, 3, 4, 5]. A very interesting and fruitful interpretation of these features is related to the description of biphoton qutrits in terms of the Stokes vectors on the Poincaré sphere [3, 4]. The qutrit state vectors (1.1) were considered as reduced to the form of two creation operators acting on the vacuum state $|\Psi\rangle = NA^\dagger B^\dagger|0\rangle$. Below, such representation of biphoton qutrits will be referred to as the “operator factorization”, and the operators $A^\dagger$ and $B^\dagger$, as “factorizing operators”. In [3, 4] the factorizing operators were considered as functions of angles determining the orientations of the corresponding Stokes vectors on the Poincaré sphere. The degree of polarization of qutrits was found to be related to the angle between these Stokes vectors. In particular, maximally polarized biphoton qutris were shown to be represented by two parallel Stokes vectors and unpolarized biphoton qutrits, by a pair of counter-directed Stokes vectors on the Poincaré sphere.

As to the entanglement of biphoton qutrits, maybe strange enough, but conclusions about its existence and degree (if it exists) were often and still remain controversial (for comparison, see, e.g., Refs. [6] and [7]). The controversy arises from definitions, formulated clearly or not. The simplest definition for pure biphoton states is well known. For biphoton qutrits this definition is formulated in terms of the wave function $\Psi(\sigma_1, \sigma_2) = \langle\sigma_1, \sigma_2|\Psi\rangle$ depending on two polarization variables of two indistinguishable photons σ_1 and σ_2. Each of these two variables can take independently one of two values $\sigma_{1,2} = H\,\text{or}\,V$. The state is entangled if the wave function cannot be factorized, $\Psi(\sigma_1, \sigma_2) \neq \varphi(\sigma_1)\chi(\sigma_2)$. In the frame of this definition and with the obligatory symmetry of biphoton wave functions taken into account, the Wootters concurrence C [8] characterizing the degree of entanglement of biphoton qutrits was found in recent papers [9, 10] to be given by

$$C = |2C_1C_3 - C_2^2|. \tag{1.2}$$

Moreover, it was found that there is a simple and direct relation between the concurrence C and the degree of polarization P for biphoton qutrits,

$$C^2 + P^2 = 1. \tag{1.3}$$

This relation shows that maximally entangled states ($C = 1$) are unpolarized and, oppositely, maximally polarized states ($P = 1$) are disentangled ($C = 0$) and their wave function is factorized. In particular, as follows from Eqs. (1.2) and (1.3), the state $|1_H, 1_V\rangle$ ($C_1 = C_3 = 0$, $C_2 = 1$) is unpolarized and maximally entangled ($P = 0$ and $C = 1$). These and other related conclusions are strongly supported by the analysis based on the Schmidt-mode decomposition [9, 10].

In this work we continue investigating the links between entanglement and polarization of biphoton qutrits, between the approaches based on the Schmidt decomposition and on the analysis of polarization Stokes vectors on the Poincaré sphere. After a brief reminder about the definitions of the Schmidt modes as the eigenfunctions of the reduced density matrix in the next section, in section 3 we suggest a very simple procedure of the operator factorization. It shows, in particular, that the operator factorization is always doable, and for any given configuration of the qutrit parameters $C_{1.2.3}$ there is only one pair of factorizing operators. In section 4 we describe an alternative procedure of the operator factorization based on the use of the Schmidt-mode representation. By combining these two approaches we show that there is a way of finding the Schmidt modes of a biphoton without solving the eigenvalue-eigenfunction problem (section 5). Finally, in section 6 we describe representations of biphoton qutrits in terms of the Stokes vectors of either the Schmidt modes or the factorizing operators. In this way we establish the full correspondence between approaches based on the Schmidt-mode and the Stokes-vector analyzes.

2. The Schmidt modes

The Schmidt modes are known to be defined as the eigenfunctions of bipartite reduced density matrices. For biphoton qutrits, the derivation is rather simple and straightforward [9, 10]. As mentioned above, the first step is finding the wave function $\Psi(\sigma_1, \sigma_2)$ corresponding to the state vector (1.1). Not reproducing here $\Psi(\sigma_1, \sigma_2)$ completely, let us show only its unfactorable part corresponding to the state $|1_H, 1_V\rangle$:

$$\Psi_{HV}(\sigma_1, \sigma_2) = \langle\sigma_1, \sigma_2|1_H, 1_V\rangle =$$
$$\frac{1}{\sqrt{2}}\Big[\delta_{\sigma_1,H}\delta_{\sigma_2,V} + \delta_{\sigma_1,V}\delta_{\sigma_2,H}\Big] \equiv \frac{1}{\sqrt{2}}\left[\binom{1}{0}_1 \otimes \binom{0}{1}_2 + \binom{0}{1}_1 \otimes \binom{1}{0}_2\right]. \tag{2.1}$$

The two alternative expressions in the second line of this equation correspond to the equivalent functional and matrix forms of writing wave functions. They both contain two terms because the wave function of two photons (bosons) has to be symmetric with respect to the transposition of variables $\sigma_1 \rightleftharpoons \sigma_2$. In the matrix form the upper and lower lines of the columns correspond to the horizontal and vertical polarizations with subscripts 1 and 2 indicating photon variables, e.g., as $\left(\begin{smallmatrix}1\\0\end{smallmatrix}\right)_1 \equiv \delta_{\sigma_1,H}$ etc.

The next step consists of constructing the density matrix $\rho = \Psi \otimes \Psi^\dagger$ (Eq. (4.2) of Ref. [9]) and its reduction, i.e., taking its trace with respect to one of the variables, e.g. variable 2. This yields [9]

$$\rho_r = \left(\begin{array}{cc} |C_1|^2 + \dfrac{|C_2|^2}{2} & \dfrac{C_1C_2^* + C_2C_3^*}{\sqrt{2}} \\ \dfrac{C_1^*C_2 + C_2^*C_3}{\sqrt{2}} & |C_3|^2 + \dfrac{|C_2|^2}{2} \end{array}\right). \tag{2.2}$$

As photons are indistinguishable particles and therefore the qutrit wave function has to be symmetric, Eq. (2.2) is also valid for the second reduced density matrix, which can be obtained by taking the trace over variable 1. The Schmidt modes are the eigenfunctions of the joint reduced density matrix (2.2),

$$\rho_r\varphi_\pm = \lambda_\pm\varphi_\pm, \tag{2.3}$$

where $\lambda_\pm$ are the eigenvalues of ρ_r related to the qutrit concurrence C (1.2) by equations

$$\lambda_\pm = \frac{1}{2}\Big(1 \pm \sqrt{1-C^2}\Big) = \frac{1 \pm P}{2} \quad \text{and} \quad C = 2\sqrt{\lambda_+\lambda_-}. \tag{2.4}$$

The Schmidt decomposition for the qutrit wave function has the form

$$\Psi(\sigma_1, \sigma_2) = \sqrt{\lambda_+}\;\varphi_+(\sigma_1)\varphi_+(\sigma_2) + \sqrt{\lambda_-}\;\varphi_-(\sigma_1)\varphi_-(\sigma_2). \tag{2.5}$$

Because of the indistinguishability of the photons, pairs of Schmidt modes (φ_+, φ_+) and (φ_-, φ_-) are formed by identical rather than different functions. This means in fact that each of the two terms in the Schmidt decomposition (2.5) describes a state with two photons in the same mode, (+) or (−) [11]. For finding the Schmidt modes explicitly, one has to solve Eq. (2.3). In the general case such solutions are rather cumbersome. An alternative approach is described below.

The polarization Stokes vector and the degree of polarization can be defined for the biphoton qutrit in terms of the reduced density matrix ρ_r as

$$\vec{S} = Tr(\rho_r\vec{\sigma}) \quad \text{and} \quad P = |\vec{S}|, \tag{2.6}$$

where now $\vec{\sigma}$ is the vector of the Pauli matrices. From here we can deduce immediately that the reduced density matrix of biphoton qutrits (2.2) coincides with the well-known polarization matrix

$$\rho_r = \rho_{pol} = \frac{1}{2}\left(\begin{array}{cc} 1+S_3 & S_1 - iS_2 \\ S_1 + iS_2 & 1 - S_3 \end{array}\right). \tag{2.7}$$

The polarization matrix is written here in its standard form [12] although with the components of the Stokes vector re-numbered. Provided that in the collinear SPDC the biphoton beam propagates along the z-axis with the horizontal and vertical directions denoted by x and y, the Stokes parameter $S_3 = 1$ corresponds to the linear polarization along the x-axis, $S_3 = -1$ corresponds to the linear polarization along the y-axis, $S_1 = \pm 1$ corresponds to the linear polarization along the lines directed at $\pm 45^\circ$ to the x-axis in the xy-plane, and $S_2 = \pm 1$, to circular polarizations.

3. Operator factorization

According to the idea of Ref. [3], the qutrit state vector (1.1) with arbitrary parameters $C_{1,2,3}$ can be reduced to the form of a product of two new creation operators times the vacuum state,

$$|\Psi\rangle = N A^\dagger B^\dagger |0\rangle, \tag{3.1}$$

where N is the normalization factor such that $\langle\Psi|\Psi\rangle = 1$. The fact that representation (3.1) always exists and is unique, already mentioned in Ref. [3], can be very easily proved as follows. Consider an operator $\hat{Q}$ such that $|\Psi\rangle = \hat{Q}|0\rangle$,

$$\hat{Q} = \frac{C_1}{\sqrt{2}}\, a_H^{\dagger 2} + C_2\, a_H^\dagger a_V^\dagger + \frac{C_3}{\sqrt{2}}\, a_V^{\dagger 2}, \tag{3.2}$$

and a second-order polynomial $Q(x)$ of some variable x, associated with $\hat{Q}$,

$$Q(x) = \frac{C_1}{\sqrt{2}}\, x^2 + C_2\, x + \frac{C_3}{\sqrt{2}}. \tag{3.3}$$

By finding the roots of the equation $Q(x) = 0$,

$$x_{A,B} = \frac{-C_2 \pm \sqrt{C_2^2 - 2C_1C_3}}{\sqrt{2}\, C_1}, \tag{3.4}$$

we represent $Q(x)$ as a product of two linear functions of x,

$$Q(x) = \frac{C_1}{\sqrt{2}}\,(x - x_A)(x - x_B). \tag{3.5}$$

As the operators $a_H^\dagger$ and $a_V^\dagger$ commute with each other, they can be arbitrarily transposed and combined as simple numbers. Hence, we can immediately write down the operator analog of the algebraic equality (3.5),

$$\hat{Q} = \frac{C_1}{\sqrt{2}}\,(a_H^\dagger - x_A\, a_V^\dagger)(a_H^\dagger - x_B a_V^\dagger). \tag{3.6}$$

This equation determines the factorizing operators $A^\dagger$ and $B^\dagger$ of Eq. (3.1), though it leaves undefined their phases φ_0 and $-\varphi_0$:

$$A^\dagger = e^{i\varphi_0}\frac{a_H^\dagger - x_A\, a_V^\dagger}{\sqrt{1+|x_A|^2}} \quad \text{and} \quad B^\dagger = e^{-i\varphi_0}\frac{a_H^\dagger - x_B a_V^\dagger}{\sqrt{1+|x_B|^2}}, \tag{3.7}$$

The phase factors in these definitions do not affect usual commutation rules for the photon annihilation and creation operators in a given mode,

$$[A, A^\dagger] \equiv AA^\dagger - A^\dagger A = 1 \text{ and } [B, B^\dagger] = 1. \tag{3.8}$$

But they do affect the cross commutators $[A, B^\dagger]$ and $[B, A^\dagger] = [A, B^\dagger]^\dagger$. As shown in the following section, these commutators are always real, whereas the parameters $x_{A,B}$ (3.4) can be complex, $x_{A,B} = |x_{A,B}|\exp(i\varphi_{A,B})$, where φ_A and φ_B are phases of x_A and x_B. Thus, the phase φ_0 can be found just from the condition that the cross-commutators are real

$$[A, B^\dagger] = \frac{e^{-2i\varphi_0}(1 + x_A^* x_B)}{\sqrt{(1 + |x_A|^2)(1 + |x_B|^2)}} = \frac{|1 + x_A^* x_B|}{\sqrt{(1 + |x_A|^2)(1 + |x_B|^2)}}. \tag{3.9}$$

Explicitly, the phase φ_0 is given by

$$\varphi_0 = \frac{1}{2}\tan^{-1}\left[\frac{|x_A||x_B|\sin(\varphi_B - \varphi_A)}{1 + |x_A||x_B|\cos(\varphi_B - \varphi_A)}\right]. \tag{3.10}$$

The commutator (3.9) is a very important characteristics of the biphoton qutrit states and, in particular, it determines the normalizing factor N in Eq. (3.1)

$$N = \frac{1}{\sqrt{1 + [A, B^\dagger]^2}}. \tag{3.11}$$

4. Operator factorization in the Schmidt representation

The Schmidt decomposition (2.5) provides an alternative way for the operator factorization of qutrit state vectors. In this approach we can further specify the earlier introduced parameters, such as the commutator of the operators A and $B^\dagger$ (3.9), as well as the definition (3.7) of the operators $A^\dagger$ and $B^\dagger$ themselves.

As mentioned above, owing to the symmetry of the qutrit wave functions, the Schmidt decomposition (2.5) presents qutrits as a coherent superposition of two orthogonal two-photon states, $|2_+\rangle$ and $|2_-\rangle$. For this reason, directly from Eq. (2.5) we can find the operator Schmidt decomposition of the qutrit state vector,

$$|\Psi\rangle = \frac{1}{\sqrt{2}}\left(\sqrt{\lambda_+}\, {a_+^\dagger}^2 + \sqrt{\lambda_-}\, {a_-^\dagger}^2\right)|0\rangle, \tag{4.1}$$

where $a_+^\dagger$ and $a_-^\dagger$ are the photon creation operators for the Schmidt modes $(+)$ and $(-)$. In terms of $a_\pm^\dagger$ the operator $\hat{Q}$ (3.2) takes the form $\hat{Q} = \left(\sqrt{\lambda_+}\, {a_+^\dagger}^2 + \sqrt{\lambda_-}\, {a_-^\dagger}^2\right)/\sqrt{2}$, and it is easily factorized to give the above-defined and appropriately normalized operators $A^\dagger$ and $B^\dagger$ in the Schmidt representation,

$$\begin{aligned} A^\dagger &= \frac{\lambda_+^{1/4} a_+^\dagger + i\lambda_-^{1/4} a_-^\dagger}{\left(\sqrt{\lambda_+} + \sqrt{\lambda_-}\right)^{1/2}} \equiv \frac{\lambda_+^{1/4} a_+^\dagger + i\lambda_-^{1/4} a_-^\dagger}{(1 + C)^{1/4}}, \\ B^\dagger &= \frac{\lambda_+^{1/4} a_+^\dagger - i\lambda_-^{1/4} a_-^\dagger}{\left(\sqrt{\lambda_+} + \sqrt{\lambda_-}\right)^{1/2}} \equiv \frac{\lambda_+^{1/4} a_+^\dagger - i\lambda_-^{1/4} a_-^\dagger}{(1 + C)^{1/4}}. \end{aligned} \tag{4.2}$$

The commutator of the operators A and $B^\dagger$ can be now written in terms of the Schmidt eigenvalues $\lambda_\pm$, or the concurrence C, or the degree of polarization P,

$$[A, B^\dagger] = [B, A^\dagger] = \frac{\sqrt{\lambda_+} - \sqrt{\lambda_-}}{\sqrt{\lambda_+} + \sqrt{\lambda_-}} = \sqrt{\frac{1 - C}{1 + C}} = \frac{P}{1 + \sqrt{1 - P^2}}, \tag{4.3}$$

and similarly for the normalization factor in Eq. (3.1),

$$N = \frac{1}{\sqrt{1+[A,B^\dagger]^2}} = \frac{\sqrt{\lambda_+}+\sqrt{\lambda_-}}{\sqrt{2}} = \sqrt{\frac{1+C}{2}}. \tag{4.4}$$

Eq. (4.3) shows that in the maximally entangled unpolarized states ($C=1$, $P=0$) the commutator $[A,B^\dagger]$ equals zero, which means that the operators $A^\dagger$ and $B^\dagger$ represent orthogonal modes, as, e.g., $a_H^\dagger$ and $a_V^\dagger$. In the case of disentangled maximally polarized states ($C=0$, $P=1$) the commutator $[A,B^\dagger]$ equals 1, which indicates that the operators $A^\dagger$ and $B^\dagger$ coincide with each other. Eq. (4.3) can be inverted to express the concurrence C via the commutator $[A,B^\dagger]$,

$$C = \frac{1-[A,B^\dagger]^2}{1+[A,B^\dagger]^2}. \tag{4.5}$$

In principle, this formula can be considered as a definition of the concurrence alternative to that of Eq. (1.2). Though it is hardly easy to find the commutator $[A,B^\dagger]$ before finding the concurrence, Eq. (4.5) is important because it shows that for getting disentangled states of biphoton qutrits it is not sufficient to provide the operator factorization. It is necessary to have the commutator equal to unity, $[A,B^\dagger]=1$. In other words, the operator factorization by no means leads automatically to the factorization of the biphoton wave function and, hence, the operator factorization does not guarantee disentanglement. The simplest and most often discussed example is the state $|1_H,1_V\rangle$. The operator factorization is present automatically with the factorizing operators $A^\dagger=a_H^\dagger$ and $B^\dagger=a_V^\dagger$. But the commutator $[A,B^\dagger]=[a_H,a_V^\dagger]$ equals zero, and hence, there is no factorization in the biphoton wave function (2.1), the state is maximally entangled and unpolarized.

5. Finding the Schmidt modes without solving the eigenvalue-eigenfunction equation (2.3)

Eqs. (4.2) can be inverted to express the Schmidt-mode creation operators $a_+^\dagger$ and $a_-^\dagger$ in terms of the factorizing operators $A^\dagger$ and $B^\dagger$

$$\begin{aligned} a_+^\dagger &= \frac{1}{2}\left(\frac{1+C}{\lambda_+}\right)^{1/4}(A^\dagger+B^\dagger),\\ a_-^\dagger &= \frac{i}{2}\left(\frac{1+C}{\lambda_-}\right)^{1/4}(B^\dagger-A^\dagger). \end{aligned} \tag{5.1}$$

By substituting here expressions (3.7) for $A^\dagger$ and $B^\dagger$ via $a_H^\dagger$ and $a_V^\dagger$, we finally get the general expressions for the Schmidt-mode creation operators,

$$\begin{aligned} a_+^\dagger = \frac{1}{2}\left(\frac{1+C}{\lambda_+}\right)^{1/4}\Bigg\{&\left[\frac{e^{i\varphi_0}}{\sqrt{1+|x_A|^2}}+\frac{e^{-i\varphi_0}}{\sqrt{1+|x_B|^2}}\right]a_H^\dagger\\ &-\left[\frac{e^{i\varphi_0}x_A}{\sqrt{1+|x_A|^2}}+\frac{e^{-i\varphi_0}x_B}{\sqrt{1+|x_B|^2}}\right]a_V^\dagger\Bigg\} \end{aligned} \tag{5.2}$$

and

$$a_-^\dagger = \frac{1}{2}\left(\frac{1+C}{\lambda_-}\right)^{1/4}\left\{\left[\frac{e^{-i\varphi_0}}{\sqrt{1+|x_B|^2}} - \frac{e^{i\varphi_0}}{\sqrt{1+|x_A|^2}}\right] a_H^\dagger \right.$$
$$\left. - \left[\frac{e^{-i\varphi_0}x_B}{\sqrt{1+|x_B|^2}} - \frac{e^{i\varphi_0}x_A}{\sqrt{1+|x_A|^2}}\right] a_V^\dagger\right\}. \tag{5.3}$$

These operators can be checked to obey the usual commutation rules for the creation and annihilation operators in two orthogonal modes,

$$[a_+, a_+^\dagger] = 1,\ [a_-, a_-^\dagger] = 1,\ [a_+, a_-^\dagger] = [a_-, a_+^\dagger] = 0. \tag{5.4}$$

The Schmidt modes corresponding to the operators $a_+^\dagger$ (5.2) and $a_-^\dagger$ (5.3) are given by

$$\varphi_\pm(\sigma) = \langle\sigma|a_\pm^\dagger|0\rangle \equiv \begin{pmatrix}\langle 1_H|a_\pm^\dagger|0\rangle \\ \langle 1_V|a_\pm^\dagger|0\rangle\end{pmatrix}. \tag{5.5}$$

It should be stressed that the Schmidt modes are found here with the help of simple algebraic operations and without solving the eigenvalue-eigenfunction equation (2.3), which is rather unexpected.

As an example, let us consider a simplified biphoton qutrit of the form

$$|\Psi\rangle = N\, a_H^\dagger\,(\cos\alpha\, a_H^\dagger + \sin\alpha\, a_V^\dagger)|0\rangle. \tag{5.6}$$

This is the case when the qutrit state vector "automatically" has the form $NA^\dagger B^\dagger|0\rangle$ and no factorization procedure is needed. The factorizing operators $A^\dagger$ and $B^\dagger$ are then given by

$$A^\dagger = a_H^\dagger \quad \text{and} \quad B^\dagger = \cos\alpha\, a_H^\dagger + \sin\alpha\, a_V^\dagger. \tag{5.7}$$

Their commutator and normalization constant N are equal to

$$[A, B^\dagger] = \cos\alpha \quad \text{and} \quad N = \frac{1}{\sqrt{1+\cos^2\alpha}}. \tag{5.8}$$

In the original form (1.1), the parameters of the qutrit (5.6) are equal to

$$C_1 = \frac{\sqrt{2}\ \cos\alpha}{\sqrt{1+\cos^2\alpha}}, \quad C_2 = \frac{\sin\alpha}{\sqrt{1+\cos^2\alpha}}, \quad C_3 = 0. \tag{5.9}$$

These parameters correspond to the following concurrence C, degree of polarization P, and the eigenvalues of the reduced density matrix $\lambda_\pm$:

$$C = C_2^2 = \frac{\sin^2\alpha}{1+\cos^2\alpha}, \quad P = \sqrt{1-C^2} = \frac{2\cos\alpha}{1+\cos^2\alpha}, \tag{5.10}$$

$$\lambda_\pm = \frac{1}{2}\left(1 \pm \sqrt{1-C^2}\right) = \frac{1\pm P}{2} = \frac{1}{2}\,\frac{(1\pm\cos\alpha)^2}{1+\cos^2\alpha}. \tag{5.11}$$

As the factorizing operators (5.7) of the qutrit (5.6) are known "automatically", the Schmidt-mode creation operators can be found directly from Eqs. (5.1) rather than from more complicated equations (5.2) and (5.3),

$$\begin{aligned} a_+^\dagger &= \frac{1}{\sqrt{2(1+\cos\alpha)}}\left[(1+\cos\alpha)\, a_H^\dagger + \sin\alpha\, a_V^\dagger\right], \\ a_-^\dagger &= \frac{i}{\sqrt{2(1-\cos\alpha)}}\left[(-1+\cos\alpha)\, a_H^\dagger + \sin\alpha\, a_V^\dagger\right]. \end{aligned} \tag{5.12}$$

From these equations and with the help of Eqs. (5.5) we easily find explicit expressions for the Schmidt modes of the biphoton qutrit (5.6)

$$\varphi_+ = \frac{1}{\sqrt{2(1+\cos\alpha)}}\begin{pmatrix} 1+\cos\alpha \\ \sin\alpha \end{pmatrix} = \begin{pmatrix} \cos(\alpha/2) \\ \sin(\alpha/2) \end{pmatrix}, \tag{5.13}$$

$$\varphi_- = \frac{i}{\sqrt{2(1-\cos\alpha)}}\begin{pmatrix} -1+\cos\alpha \\ \sin\alpha \end{pmatrix} = i\begin{pmatrix} -\sin(\alpha/2) \\ \cos(\alpha/2) \end{pmatrix}. \tag{5.14}$$

6. Poincaré-sphere representation

6.1. The Stokes vectors of single-photon states

Following earlier works [3, 4] let us consider representation of the above-derived results in the space of Stokes vectors on the Poincaré-sphere. Let us start with the general expressions (3.7) for the factorizing operators $A^\dagger$ and $B^\dagger$. These operators generate a pair of single-photon states and the corresponding wave functions:

$$|1_A\rangle = A^\dagger|0\rangle \equiv \frac{1}{\sqrt{1+|x_A|^2}}(a_H^\dagger - x_A a_V^\dagger)|0\rangle, \tag{6.1}$$

$$|1_B\rangle = B^\dagger|0\rangle \equiv \frac{1}{\sqrt{1+|x_B|^2}}(a_H^\dagger - x_B a_V^\dagger)|0\rangle \tag{6.2}$$

and

$$\psi_A = \frac{1}{\sqrt{1+|x_A|^2}}\begin{pmatrix} 1 \\ -x_A \end{pmatrix},\ \psi_B = \frac{1}{\sqrt{1+|x_B|^2}}\begin{pmatrix} 1 \\ -x_B \end{pmatrix}, \tag{6.3}$$

where, as shown above, the parameters $x_{A,B}$ can be expressed via the original qutrit parameters by equations (3.4). We have dropped the phase factors occurring in operators $A^\dagger$ and $B^\dagger$ (3.7) because they do not affect the polarization matrices of the states considered below.

The density matrices of the single-photon states $|1_A\rangle$ and $|1_B\rangle$ are easily found from their wave functions to be given by

$$\rho_A = \frac{1}{1+|x_A|^2}\begin{pmatrix} 1 & -x_A^* \\ -x_A & |x_A|^2 \end{pmatrix}, \tag{6.4}$$

and the same for ρ_B with the substitution $x_A \to x_B$. By identifying the density matrices of the states $|1_A\rangle$ and $|1_B\rangle$ with the corresponding polarization matrices $\rho_{pol}^{(A)}$ and $\rho_{pol}^{(B)}$ (having the same form as the polarization matrix in Eq. (2.7)) we can find the components of the Stokes vectors $\vec{S}^{(A)}$ and $\vec{S}^{(B)}$ of the states $|1_A\rangle$ and $|1_B\rangle$:

$$\begin{aligned} S_3^{(A)} &= (1-|x_A|^2)/(1+|x_A|^2) \equiv \cos\theta_A, \\ S_1^{(A)} &= -2\mathrm{Re}(x_A)/(1+|x_A|^2) \equiv \sin\theta_A\,\cos\varphi_A, \\ S_2^{(A)} &= -2\mathrm{Im}(x_A)/(1+|x_A|^2) \equiv \sin\theta_A\,\sin\varphi_A, \end{aligned} \tag{6.5}$$

$$\begin{aligned} S_3^{(B)} &= (1-|x_B|^2)/(1+|x_B|^2) \equiv \cos\theta_B, \\ S_1^{(B)} &= -2\mathrm{Re}(x_B)/(1+|x_B|^2) \equiv \sin\theta_B\,\cos\varphi_B, \\ S_2^{(B)} &= -2\mathrm{Im}(x_B)/(1+|x_B|^2) \equiv \sin\theta_B\,\sin\varphi_B, \end{aligned} \tag{6.6}$$

where θ_A and θ_B are the angles between the vectors $\vec{S}^{\,(A)}$ and $\vec{S}^{\,(B)}$ and the x-axis on the Poincaré sphere (Fig.1). The values φ_A and φ_B are the angles between the horizontal axis $O45^\circ \perp Ox$ on the Poincaré sphere and projections of the Stokes vectors $\vec{S}^{\,(A)}$ and $\vec{S}^{\,(B)}$ on the plane perpendicular to Ox.



Figure 1. The Stokes vectors $\vec{S}^{(A)}$ (with spherical coordinates θ_A, φ_A) and $\vec{S}^{(B)}$ (6.5,6.6), as well as the biphoton Stokes vector $\vec{S}^{\,biph}$ (6.11), (6.19), (6.20), on the Poincaré sphere.

As it follows from Eqs. (6.5) and (6.6), the absolute values of the Stokes vectors $\vec{S}^{\,(A)}$ and $\vec{S}^{\,(B)}$ are equal to unity, $\left|\vec{S}^{\,(A)}\right| = \left|\vec{S}^{\,(B)}\right| = 1$. The angle between these vectors θ_{AB} is determined by their scalar product,

$$\cos\theta_{AB} = \vec{S}^{\,(A)}\cdot\vec{S}^{\,(B)} = \cos\theta_A\cos\theta_B + \sin\theta_A\sin\theta_B\cos(\varphi_A-\varphi_B). \tag{6.7}$$

6.2. Schmidt-mode representation and Poincaré-sphere images of biphoton qutrits

Let us analyze now the Poincaré-sphere image of the general-form biphton qutrit (3.1). One possibility of its investigation consists of finding from this equation the original qutrit's parameters $C_{1,2,3}$, substituting them into the general expression for the reduced density matrix (2.7), and finding from this last equation the biphoton Stokes parameters, which involves rather cumbersome transformations and formulas. An alternative, more elegant and even more informative approach is based on the use of the Schmidt-mode representation. In this representation the Schmidt modes and their single-photon density matrices are very simple:

$$\varphi_+ = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \varphi_- = \begin{pmatrix} 0 \\ 1 \end{pmatrix}; \quad \rho_+ = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \rho_- = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}. \tag{6.8}$$

The Stokes vectors $\vec{S}_+$ and $\vec{S}_-$ corresponding to the Schmidt modes (6.8) have components

$$S_3^{(+)} = 1,\ S_2^{(+)} = S_1^{(+)} = 0 \quad \text{and} \quad S_3^{(-)} = -1,\ S_2^{(-)} = S_1^{(-)} = 0. \tag{6.9}$$

In terms of the Poincaré sphere this means that the transition to the Schmidt-mode representation is realized by means of a rotation of the Poincaré sphere that makes the Stokes vectors of the Schmidt modes directed along the former (H, V) axis. Then this axis turns into the $(+,-)$ axis, so that the Stokes vectors $\vec{S}^{(+)}$ and $\vec{S}^{(-)}$ point, correspondingly, at the positive and negative directions along this axis (Fig. 2). The biphoton reduced density matrix (2.7) is also very simple in the Schmidt-mode representation


$\frac{(+)+i(-)}{\sqrt{2}}$
$\vec{S}^{(A)}$
$(-)$
$\vec{S}^{(-)}$
$\frac{(+)-(-)}{\sqrt{2}}$
$\vec{S}^{(+)}$
$\vec{S}^{biph}$
$\frac{(+)+(-)}{\sqrt{2}}$
$(+)$
$\vec{S}^{(B)}$
$\frac{(+)-i(-)}{\sqrt{2}}$
a
$\frac{(+)+i(-)}{\sqrt{2}}$
$(-)$
$\lambda_- 2\vec{S}^{(-)}$
$\frac{(+)-(-)}{\sqrt{2}}$
$\lambda_+ 2\vec{S}^{(+)}$
$\vec{S}^{biph}$
$\frac{(+)+(-)}{\sqrt{2}}$
$(+)$
$\frac{(+)-i(-)}{\sqrt{2}}$
b

Figure 2. *a*: Poincaré sphere and the Stokes vectors in the Schmidt-mode representation; *b*: A scheme of two Schmidt-mode two-photon states

$$\rho_r = \begin{pmatrix} \lambda_+ & 0 \\ 0 & \lambda_- \end{pmatrix}. \tag{6.10}$$

As well as the one-photon Schmidt-mode Stokes vectors $\vec{S}^{(\pm)}$, the biphoton Stokes vector $\vec{S}^{\,biph}$ has only one non-zero component,

$$S_3^{biph} = \lambda_+ - \lambda_- = P \quad \text{with} \quad S_2^{biph} = S_1^{biph} = 0. \tag{6.11}$$

On the Poincaré sphere the vector $\vec{S}^{\,biph}$ is directed along the same axis $(+,-)$ as the Schmidt-mode Stokes vectors $\vec{S}^{(\pm)}$. As usual, the absolute value of $\vec{S}^{\,biph}$ coincides with the degree of polarization per photon P for biphoton polarization qutrits [3, 4],

$$\left|\vec{S}^{\,biph}\right| = S_3^{biph} = P. \tag{6.12}$$

Finally, the factorizing operators $A^\dagger$ and $B^\dagger$ are given by superpositions of the Schmidt-mode creation operators (4.2). In the Schmidt-mode representation the one-photon states generated by these operators, $A^\dagger|0\rangle$ and$B^\dagger|0\rangle$, have the following wave functions

$$\psi_A = \frac{1}{(1+C)^{1/4}}\begin{pmatrix} \lambda_+^{1/4} \\ i\lambda_-^{1/4} \end{pmatrix}, \quad \psi_B = \frac{1}{(1+C)^{1/4}}\begin{pmatrix} \lambda_+^{1/4} \\ -i\lambda_-^{1/4} \end{pmatrix} \tag{6.13}$$

and density matrices

$$\rho_{A,B} = \frac{1}{\sqrt{1+C}}\begin{pmatrix} \sqrt{\lambda_+} & \mp i(\lambda_+\lambda_-)^{1/4} \\ \pm i(\lambda_+\lambda_-)^{1/4} & \sqrt{\lambda_-} \end{pmatrix}. \tag{6.14}$$

Components of the Stokes vectors of these states are $S_1^{(A)} = S_1^{(B)} = 0$ and

$$\begin{aligned} S_3^{(A)} = S_3^{(B)} &= \frac{\sqrt{\lambda_+}-\sqrt{\lambda_-}}{\sqrt{1+C}} = \sqrt{\frac{1-C}{1+C}}, \\ S_2^{(A)} = -S_2^{(B)} &= \frac{2(\lambda_+\lambda_-)^{1/4}}{\sqrt{1+C}} = \sqrt{\frac{2C}{1+C}}. \end{aligned} \tag{6.15}$$

Thus, in the Schmidt-mode representation the Stokes vectors $\vec{S}^{(A)}$ and $\vec{S}^{(B)}$ are located in the vertical plane containing the horizontal axis $(+,-)$, and their directions are symmetric with respect to this axis. The angles between $\vec{S}^{(A,B)}$ and the biphoton Stokes vector $\vec{S}^{\,biph}\|(+,-)$ are determined by $S_3^{(A,B)}$-projections of the vectors $\vec{S}^{(A,B)}$ on the $(+,-)$:

$$\cos\left(\theta_{AB}/2\right) = S_3^{(A)} = S_3^{(B)} = \sqrt{\frac{1-C}{1+C}} = \frac{P}{1+\sqrt{1-P^2}} = [A, B^\dagger], \tag{6.16}$$

where θ_{AB} is the angle between the vectors $\vec{S}^{(A)}$ and $\vec{S}^{(B)}$.

By inverting the relation between $\cos\left(\theta_{AB}/2\right)$ and the concurrence C, we can express the latter in terms of the angle θ_{AB} between the vectors $\vec{S}^{(A)}$ and $\vec{S}^{(B)}$:

$$C = \frac{1-\cos^2(\theta_{AB}/2)}{1+\cos^2(\theta_{AB}/2)} = \frac{1-\cos\theta_{AB}}{3+\cos\theta_{AB}}. \tag{6.17}$$

And the expression for the degree of polarization P via θ_{AB} has the form

$$P = \sqrt{1-C^2} = \frac{2\sqrt{2(1+\cos\theta_{AB})}}{3+\cos\theta_{AB}} = \frac{4|\cos(\theta_{AB}/2)|}{3+\cos\theta_{AB}}. \tag{6.18}$$

This result for the degree of polarization agrees perfectly with that of the paper [4].

As the biphoton Stokes vector is directed along the bisector of the angle between the Stokes vectors $\vec{S}^{(A)}$ and $\vec{S}^{(B)}$, in the vectorial form the biphoton Stokes vector can be written as

$$\begin{aligned} \vec{S}^{\,biph} &= \left|\vec{S}^{\,biph}\right| \frac{\vec{S}^{(A)}+\vec{S}^{(B)}}{\left|\vec{S}^{(A)}+\vec{S}^{(B)}\right|} = P\frac{\vec{S}^{(A)}+\vec{S}^{(B)}}{2(1+\cos\theta_{AB})} \\ &= \frac{2\cos(\theta_{AB}/2)}{(3+\cos\theta_{AB})(1+\cos\theta_{AB})}\left(\vec{S}^{(A)}+\vec{S}^{(B)}\right). \end{aligned} \tag{6.19}$$

Figure 2*b* illustrates an alternative representation of a qutrit in terms of the Schmidt-mode Stokes vectors $\vec{S}^{(+)}$ and $\vec{S}^{(-)}$. The picture shows pairs of these Stokes vectors, oppositely directed and having the weight coefficients λ_+ and λ_-. The total biphoton Stokes vector equals the sum of all four Stokes vectors of the Schmidt modes with the weight coefficients, and half of this sum gives biphoton Stokes vector per one photon:

$$\vec{S}^{\,biph} = \lambda_+ \vec{S}^{(+)} + \lambda_- \vec{S}^{(-)}\,. \tag{6.20}$$

The results of this section are invariant with respect to rotations of the Poincaré sphere. The only changes concern the orientation of the $(+)-(-)$ axis and the plane containing the Stokes vectors $\vec{S}_A$ and $\vec{S}_B$. But at any orientation of the $(+)-(-)$ axis and $\left(\vec{S}_A, \vec{S}_B\right)$-plane, the Stokes vectors of the Schmidt modes $\vec{S}^{(\pm)}$, of the factorizing operators $\vec{S}^{(A,B)}$, and of the biphoton state as a whole $\vec{S}^{\,biph}$ belong to the same plane. The biphoton Stokes vector is directed along the same direction as $\vec{S}^{(\pm)}$, and this direction coincides with the bisector of the angle θ_{AB} between $\vec{S}_A$ and $\vec{S}_B$. Eqs. (6.17)-(6.20) remain valid without any changes in any frames with rotated Poincarè sphere. As to other derived equations (6.9), (6.11), (6.12), (6.15), and (6.16), they remain valid too, though the components of Stokes vectors S_3 and S_2 have to be understood as projections on the directions of the turned $(+)-(-)$ axis and turned $\left(\vec{S}_A, \vec{S}_B\right)$-plane, rather than, e.g., projections on the $V-H$ and $R-L$ axes of the standard Poincaré-sphere orientation of Fig. 1.

7. Possible experiments

A possible experiment on the selection of polarization Schmidt modes of a qutrit is shown in Fig. 3a. The qutrit state is sent to a polarization beam splitter (PBS) preceded by a quarter-wave and a half-wave plates. The plates are oriented in such a way that photons in the polarization Schmidt mode $|\varphi_+\rangle$ become horizontally polarized [13]. Then the orthogonally polarized photons in the $|\varphi_-\rangle$ mode automatically become vertically polarized. The qutrit state vector after the plates is transformed into a weighted superposition of photon pairs in vertical and horizontal modes,

$$|\Psi\rangle = \sqrt{\lambda_+}|2_H\rangle + e^{2i\phi}\sqrt{\lambda_-}|2_V\rangle, \tag{7.1}$$

where ϕ is the relative phase of the Schmidt modes $|\varphi_-\rangle$ and $|\varphi_+\rangle$ [11].

After the PBS, the pair $|2_H\rangle$ goes into the transmitted output port and the pair $|2_V\rangle$, into the reflected output port. Each photon pair can be detected as a coincidence of single-photon detector 'clicks'. For registering such coincidences, beamsplitters (BS) followed by pairs of detectors are introduced in both output ports of the PBS. Then, coincidences will be observed between the counts of either detectors D_{1+}, D_{2+} or detectors D_{1-}, D_{2-}, and their rates $R^c_\pm$ will scale as the Schmidt eigenvalues $\lambda_\pm = R^c_\pm/(R^c_+ + R^c_-)$. But no coincidences between the outputs of the detectors in different ports of the PBS will be observed. The polarization Schmidt modes can be found experimentally by finding positions of the quarter-wave and half-wave plates at which the coincidence counting rates between the detectors in different output ports of the PBS turn to zero. This approach is most appropriate if the qutrit parameters $C_{1,2,3}$ are not known in advance. Further, from the experimentally found parameters $\lambda_\pm$, with the help of Eqs. (2.4) one can easily calculate both the qutrit concurrence and degree of polarization.

In the absence of detectors' dark counts and parasite light, the beamsplitters in the PBS output ports are not necessary and only a single detector can be used in each port. Note that this method of finding $\lambda_\pm$ can be made invariant to the quantum efficiencies of the detectors. Indeed, instead of comparing the counting (coincidence) rates at the two output ports of the PBS one can compare the counting (coincidence) rates in the same port but at two different positions of the plates, corresponding to the Schmidt mode $|\varphi_+\rangle$ or $|\varphi_-\rangle$ sent to the same port.

It is worth mentioning that the four parameters determining a qutrit can be also introduced as the orientations of the plates transforming the qutrit from the form (4.1) into the one (7.1), one of the Schmidt eigenvalues, and the phase ϕ. Experimentally, the latter can be measured in the same scheme as shown in Fig. 3a but with the PBS turned for 45°. Indeed, if the number of qutrits at the input of the PBS is N, and the quantum efficiencies of detectors D_{1+} and D_{2+} are, respectively, η_1 and η_2, then the numbers of their coincidences with the PBS oriented at 0° and 90° will be $R_0 = \eta_1\eta_2\lambda_+ N/2$ and $R_{90} = \eta_1\eta_2\lambda_- N/2$, respectively. At the same time, for a 45° orientation of the PBS the number of coincidences will be $R_{45} = \eta_1\eta_2(1 + 2\sqrt{\lambda_+\lambda_-}\cos 2\phi)N/4$. From the measurement of these three parameters, one can infer both the Schmidt eigenvalues and the phase ϕ without knowing the quantum efficiencies and the initial number of pairs:

$$\lambda_+ = \frac{R_0}{R_0 + R_{90}}, \lambda_- = \frac{R_{90}}{R_0 + R_{90}}, \cos 2\phi = \frac{2R_{45} - R_0 - R_{90}}{2\sqrt{R_0 R_{90}}}. \qquad (7.2)$$



Figure 3. *a*: Experimental setup for observing polarization Schmidt modes of a qutrit $|\Psi\rangle$; *b*: Experimental setup for demonstrating polarization entanglement of a qutrit $|\Psi\rangle$.

For the physical interpretation of the polarization entanglement of a qutrit, one can suggest another experimental scheme (Fig. 3b). The qutrit $|\Psi\rangle$, after the polarization transformation, is directed to a 50% non-polarizing beam splitter (NPBS). Half of the photon pairs will exit into a single output port of the NPBS, but the other half will be split between the two ports. This part of the state can be written as

$$|\Psi\rangle_{12} \equiv \sqrt{\lambda_+}\,|1_{H1}, 1_{H2}\rangle + e^{2i\phi}\sqrt{\lambda_-}\,|1_{V1}, 1_{V2}\rangle, \tag{7.3}$$

where the indices $H1$ ($H2$) denote horizontal polarization in the transmitted (reflected) beam, and similarly for the vertical polarization.

This state has been considered in Ref. [14], and it was shown to be maximally entangled only if the Schmidt eigenvalues are equal, $\lambda_+ = \lambda_-$. This exactly corresponds to the case where $C = 1$ and the qutrit $|\Psi\rangle$ is fully entangled. If $\lambda_+ \neq \lambda_-$, the state is non-maximally entangled, and its degree of entanglement can be measured, as shown in Ref. [14], from the orientations of the polarizers P_1, P_2 at which the coincidence counting rate for detectors D_1, D_2 vanishes.

8. Conclusion

We have considered the general case of a biphoton qutrit, physically represented by a photon pair in a single frequency and wavevector mode. We have found a simple procedure of factorizing the qutrit state vectors, i.e., representing them as products of two single-photon creation operators in certain polarization modes acting on a vacuum. The procedure clearly proves the existence and uniqueness of such a representation. We have shown that the described operator factorization of state vectors is not related to the factorization of the biphoton polarization wave function and does not exclude a possibility of entanglement for biphoton qutrits. The degrees of their entanglement and polarization are found to be related directly to the commutation features of the operators factorizing the state vectors by Eqs. (4.3) and (4.5). Further, we considered the polarization Schmidt decomposition of a generic biphoton qutrit and derived it both using the standard method and the factorization procedure. For the qutrit and its Schmidt modes, the Stokes vectors have been calculated and shown on the Poincaré sphere. Finally, an experiment has been suggested to find the polarization Schmidt modes of a qutrit and to measure its degree of entanglement and polarization. A physical interpretation is proposed for the polarization entanglement of a biphoton qutrit.

Acknowledgments

This work was supported in part by the Russian Foundation for Basic Research, grant #11-02-01074.

[1] A.V. Burlakov and D.N. Klyshko, *JETP Lett.*, **69**, 839 (1999)
[2] A.V. Burlakov, M. V. Chekhova, O. A. Karabutova, D. N. Klyshko, and S. P. Kulik *Phys. Rev.* A **60**, R4209 (1999)
[3] A.V. Burlakov, M.V. Chekhova, *JETP Lett.*, **75**, 432 (2002)
[4] A.A. Zhukov, G.A. Maslennikov, M.V. Chekhova, *JETP Lett.*, **76**, 596, (2002)

[5] M. V. Chekhova, L. A. Krivitsky, S. P. Kulik, and G. A. Maslennikov, *Phys. Rev.* A **70**, 053801 (2004)

[6] M.H. Rubin, D.N. Klyshko, Y.H. Shih, and A.V. Sergienko, *Phys. Rev.* A **50** 5122 (1994)

[7] B.P. Lanyon et al, Phys. Rev. Lett. **100** 060504 (2008)

[8] W.K. Wootters *Phys. Rev. Lett.*, **80**, 2245 (1998)

[9] M.V. Fedorov, P.A. Volkov, Yu.M. Mikhailova, S.S. Straupe, S.P. Kulik, *New J. Phys.*, **13**, 083004 (2011)

[10] M.V. Fedorov, P.A. Volkov, Yu.M. Mikhailova, *JETP*, **115**, 15 (2012)

[11] Despite the fact that the polarization state of a single photon is usually considered with the total phase meaningless and therefore set equal to zero, the relative phase ϕ between the Schmidt modes φ_- and φ_+ is important and cannot be omitted.

[12] L.D. Landau and E.M. Lifshitz, The Classical Theory of Fields, Butterworth-Heinemann Ltd, Oxford, 1975

[13] This is always possible as such a combination of plates suffices for transforming any polarization state of light into the linear polarization state transmitted by the PBS.

[14] A. G. White, D. F. V. James, P. H. Eberhard, and P. G. Kwiat, *Phys. Rev. Lett.* **83** 3103 (1999)